

QUENCHED CHIRAL PERTURBATION THEORY FOR BARYONS

James N. Labrenz¹ and Stephen R. Sharpe²

*Physics Department, FM-15
University of Washington
Seattle, WA 98195*

Abstract

We develop quenched chiral perturbation theory for baryons using the graded-symmetry formalism of Bernard and Golterman and calculate non-analytic contributions to the baryon masses coming from quenched chiral loops. The usual term proportional to $m_q^{3/2}$ is substantially altered due to the cancellation of diagrams with internal quark loops. In addition, the η' “hairpin” vertex leads to a new correction, proportional to $m_q^{1/2}$. We compare our results to numerical lattice data and use them to estimate the size of the quenching error in the octet baryon masses.

*Presented by J. Labrenz at LATTICE 93,
International Symposium on Lattice Field Theory,
Dallas, Texas, U.S.A. 12-17 October, 1993.
An abridged version will appear in the proceedings.*

December 1993

¹email: sharpe@galileo.phys.washington.edu

²email: labrenz@galileo.phys.washington.edu

1 Introduction and Formalism

There have been several recent calculations of non-analytic loop corrections, or chiral logs, for the masses and amplitudes of quenched pseudo-Goldstone bosons [1, 2]. The aim has been to study the effect of the quenched approximation (QA) as the chiral limit is approached. In this note, we extend this work to baryons, using the Lagrangian approach of Bernard and Golterman [1].

The key observation of Ref. [1] is that quenched QCD possesses a larger chiral symmetry than the full theory. Following Morel, the fermionic determinant is removed from the functional integral by adding three pseudo-fermionic (*i.e.* commuting, spin-1/2) ghost quarks $\tilde{u}$, $\tilde{d}$ and $\tilde{s}$, degenerate in mass with the physical quarks and coupled to the gauge fields in the same way. The resulting Lagrangian is symmetric, in the chiral limit, under the graded chiral symmetry $SU(3|3)_L \times SU(3|3)_R$, and this symmetry must be used to construct a low-energy chiral Lagrangian.

We introduce baryons using the “heavy-quark” formalism employed by Jenkins and Manohar in ordinary chiral perturbation theory (ChPT) [3]. Our field $\mathcal{B}_v$ satisfies all the kinematic properties of the baryon field B_v of ref. [3]—*i.e.* $\mathcal{B}_v(x) = \exp(im_{\mathcal{B}}\not{v}x^\mu)\mathcal{B}(x)$, $\mathcal{B}_v = \frac{1}{2}(1 + \not{v})\mathcal{B}_v$, and $S_v^2\mathcal{B}_v = -\frac{3}{4}\mathcal{B}_v$, where v_μ is the 4-velocity, and S_v^μ , the covariant “spin” operator, satisfies $v \cdot S_v = 0$. Henceforth we drop the subscript v .

We find that, in the QA, the field $\mathcal{B}$ must be a three index tensor, $\mathcal{B}_{ijk}$, as opposed to the usual 3×3 matrix B . In an ordinary chiral theory, $B \mapsto UBU^\dagger$, where U is introduced by defining $\xi(x) = \exp(i\pi(x)/f) = \Sigma^{1/2}(x)$, such that $\xi \mapsto L\xi U^\dagger = U\xi R^\dagger$. The latter two relations are “quenched” by replacing π with the graded field Φ , defined by eq. (3) of ref. [1]; then $L, R \in SU(3|3)_{L,R}$. To construct the baryons, we introduce the “Quark” vector $Q = (u, d, s, \tilde{u}, \tilde{d}, \tilde{s})$ which transforms under $SU(3|3)_V$ (*i.e.* $L=R=U$) as $Q_i \mapsto U_{ij}Q_j$. The baryon field is then defined to transform as a direct product of Quark vectors:

$$\mathcal{B}_{ijk}^\gamma \sim \left[Q_i^{\alpha,a} Q_j^{\beta,b} Q_k^{\gamma,c} - Q_i^{\alpha,a} Q_j^{\gamma,c} Q_k^{\beta,b} \right] \varepsilon_{abc} (C\gamma_5)_{\alpha\beta}, \quad (1)$$

where $C = i\gamma_2\gamma_0$ is the charge conjugation matrix. Generalizing to chiral transformations $U(L, R, \Phi)$, we obtain

$$\mathcal{B}_{ijk}^\gamma \mapsto (-1)^{i'(j+j')+(i'+j')(k+k')} U_{ii'} U_{jj'} U_{kk'} \mathcal{B}_{i'j'k'}^\gamma, \quad (2)$$

where we use a standard notation for the grading factor. (Indices corresponding to physical quarks (1–3) are read as 1, and those corresponding to ghost quarks (4–6) are read as 0.)

The relevant part of the quenched chiral Lagrangian including baryons is $\mathcal{L}^{(Q)} = \mathcal{L}_{BG} + \mathcal{L}_{B\Phi}$, where

$$\mathcal{L}_{BG} = \frac{f^2}{4} \left[\text{str}(\partial_\mu \Sigma \partial^\mu \Sigma^\dagger) + 2\mu \text{str}(\mathcal{M}^+) \right] + \alpha_0 \partial_\mu \Phi_0 \partial^\mu \Phi_0 - m_0^2 \Phi_0^2 \quad (3)$$

is the Bernard-Golterman Lagrangian, and

$$\begin{aligned} \mathcal{L}_{B\Phi} = & i(\overline{\mathcal{B}}v \cdot \mathcal{D}\mathcal{B}) + 2\alpha(\overline{\mathcal{B}}S^\mu \mathcal{B}A_\mu) + 2\beta(\overline{\mathcal{B}}S^\mu A_\mu \mathcal{B}) + 2\gamma(\overline{\mathcal{B}}S^\mu \mathcal{B})\text{str}(A_\mu) \\ & + \alpha_M(\overline{\mathcal{B}}\mathcal{B}\mathcal{M}^+) + \beta_M(\overline{\mathcal{B}}\mathcal{M}^+ \mathcal{B}) + \sigma(\overline{\mathcal{B}}\mathcal{B})\text{str}(\mathcal{M}^+). \end{aligned} \quad (4)$$

Here, str is the supertrace, and the additional fields are defined as

$$\begin{aligned}\Phi_0 &= \frac{1}{\sqrt{3}}\text{str}\Phi \equiv \frac{1}{\sqrt{2}}(\eta' - \tilde{\eta}'), \\ A_\mu &= i\frac{1}{2}(\xi\partial_\mu\xi^\dagger - \xi^\dagger\partial_\mu\xi), \\ \mathcal{M}^+ &= \xi^\dagger M\xi^\dagger + \xi M\xi,\end{aligned}$$

where $M = \text{diag}(m_u, m_d, m_s, m_u, m_d, m_s)$. Our convention is such that $f_\pi = 93\text{MeV}$. In $\mathcal{L}_{\mathcal{B}\Phi}$ the brackets $()$ denote group-invariant contractions. (Anti-)symmetry under $j \leftrightarrow k$ in $\mathcal{B}_{ijk}$ leaves only two unique couplings to fields such as A_i^j , and these are denoted by the ordering.

In the QA the η' (and $\tilde{\eta}'$) remain light, and the propagator is to be computed only from the Σ terms in $\mathcal{L}_{BG}$. Terms quadratic in the singlet Φ_0 are treated as additional (so-called hairpin) vertices; the insertion of the one proportional to $\delta \equiv m_0^2/(48\pi^2 f^2)$ leads to enhanced chiral logs.

One subtlety in doing calculations is that there is a several-to-one mapping between the $\mathcal{B}_{ijk}$ and the independent fields. In the physical (quark) sector $\mathcal{B}_{ijk}|_q$, where indices run from 1 to 3, we find

$$\mathcal{B}_{ijk}|_q = \frac{1}{\sqrt{6}}(\varepsilon_{ijk'}B_k^{k'} + \varepsilon_{ikk'}B_j^{k'}). \quad (5)$$

The Clebsch for a meson loop i on baryon line b can be written as

$$c_{b,i} = \sum_{\alpha,\beta} \psi_\alpha^b D_{\alpha\beta}^{b,i} \psi_\beta^b. \quad (6)$$

The ψ^b are normalization factors which can be obtained from (5). For example, for the proton,

$$\psi^p = (\psi_{112}, \psi_{121}, \psi_{211}) = \sqrt{\frac{1}{6}}(1, 1, -2). \quad (7)$$

The $D^{b,i}$ are numbers computed from the “flavor” part of the Feynman rules and can be represented as a sum of quark-flow diagrams, as illustrated below.

2 Mass Renormalization

The important loop corrections to the baryon masses m_b are shown in Fig. 1. The axial vertex ($\bullet$) is a sum of three terms, “ α ,” “ β ” and “ γ ,” corresponding to the couplings in $\mathcal{L}_{\mathcal{B}\Phi}$, and there is a similar sum for the symmetry-breaking mass vertex, denoted by the solid triangle. The baryon propagator can be written as

$$\langle B_{lmn} \overline{B}^{kji} \rangle = \frac{i}{(v \cdot k) + i\epsilon} \left[\begin{array}{c} l \ m \ n \\ | \ | \ | \\ i \ j \ k \end{array} + \begin{array}{c} l \ m \ n \\ | \ \diagdown \ \diagup \\ i \ j \ k \end{array} + \begin{array}{c} l \ m \ n \\ \diagup \ \diagdown \ | \\ i \ j \ k \end{array} + \begin{array}{c} l \ m \ n \\ \diagdown \ \diagup \ \diagdown \\ i \ j \ k \end{array} + \begin{array}{c} l \ m \ n \\ \diagup \ \diagdown \ \diagup \\ i \ j \ k \end{array} + \begin{array}{c} l \ m \ n \\ \diagup \ \diagdown \ \diagdown \\ i \ j \ k \end{array} \right],$$

where the flavor diagrams denote delta functions in the matched indices and carry Clebsch and grading factors which we have suppressed. The pion propagator is diagonal in the $Q_i \overline{Q}_j$ basis of states. In the neutral sector, this is different from ordinary ChPT, where instead the propagator is diagonal in the π^0, η, η' basis. This difference is a manifestation of how the graded symmetry “quenches” the η' and forces the introduction of the hairpin vertex.




Figure 1: Baryon mass renormalization in quenched ChPT. $\bullet$ is the axial vertex (α, β, γ) , the solid triangle is the mass vertex $(\alpha_M, \beta_M, \sigma)$, and $\times$ is the hairpin vertex $(m_0^2 = \delta 48\pi^2 f^2)$.

Consider the term from Fig. 1a proportional to β^2 . It can be written as

$$-i\Delta m_b = i\beta^2 \sum_{i=\pi, K, s\bar{s}} c_{b,i} X(M_i), \quad (8)$$

where $X(M) = M^3/(32\pi f^2)$ comes from the Feynman integral, and the Clebsch factor can be computed in terms of quark flow diagrams, as described above. For instance, if b denotes the proton, then the factors which enter into Eq. (6) are of the form

$$D_{112,112}^{p,\pi} = \left(\sum_{l=\{1,2,4,5\}} \left(\begin{array}{c} \text{diagram with } l \text{ loops} \\ 1 \ 1 \ 2 \end{array} \right) \right) + \left(\begin{array}{c} \text{diagram with } 1 \text{ loop} \\ 1 \ 1 \ 2 \end{array} \right) + \left(\begin{array}{c} \text{diagram with } 2 \text{ loops} \\ 1 \ 1 \ 2 \end{array} \right).$$

As expected, the graded symmetry produces a cancellation of all “quark loop” diagrams (the first term above is given by $\frac{1}{6} \sum_{l=\{1,1,0,0\}} (-1)^{l+1}$, where the grading arises from the pion propagator), but there remain non-zero contributions from diagrams with only “valence quark” lines. Thus the mass renormalization proportional to $M_\pi^3 \sim m_q^{3/2}$ is substantially altered from its value in ordinary ChPT. The quenched nucleon, for example, will not be renormalized by kaon loops.

Next, consider the hairpin insertion of Fig. 1b. This Feynman integral contains an additional pion propagator which reduces the order of the finite part by a power of M_π^2 . Thus a new non-analytic term (one proportional to $M_\pi \sim m_q^{1/2}$) appears in the quenched expansion for baryon mass renormalization. The flavor factor in this case receives non-vanishing contributions only from the α and β vertices; terms containing a γ vertex always have an internal quark loop. For instance,

$$\beta^2 \sim \left(\begin{array}{c} \text{diagram} \\ \text{---} \end{array} \right), \quad \alpha\beta \sim \left(\begin{array}{c} \text{diagram} \\ \text{---} \end{array} \right), \quad \beta\gamma \sim \left(\begin{array}{c} \text{diagram} \\ \text{---} \end{array} \right).$$

We have calculated the renormalization for the octet baryons up to $\mathcal{O}(M_\pi^3)$ for $m_u = m_d \neq m_s$. We first present the result for degenerate quarks. To facilitate the comparison with ordinary ChPT, we insert (5) into $\mathcal{L}_{B\Phi}$ and compute the correspondence between our parameters α, β, α_M , etc. and the canonical ones defined in [4]. We find $\alpha = 2(\frac{1}{3}D+F)$, $\beta = (-\frac{5}{3}D+F)$, $\alpha_M = 4(\frac{1}{3}b_D+b_F)$, $\beta_M = 2(-\frac{5}{3}b_D+b_F)$ and $\sigma = 2(b_0+b_D-b_F)$. The γ -vertex is unique to the quenched theory. Our result for the baryon mass is

$$m_N = \bar{m} - \frac{3\pi}{2}(D-3F)^2 \delta M_\pi + 2(b_D-3b_F) \hat{M}_\pi^2 + \left[\frac{2}{3}(D-3F)(2D+3\gamma) + \frac{5}{6}\alpha_0(D-3F)^2 \right] \frac{M_\pi^3}{16\pi f^2}, \quad (9)$$

where $M_\pi^2 = 2\mu m_q$. There is a logarithm from Fig. 1c which is absorbed into the tree-level result through pion mass renormalization:

$$\hat{M}_\pi^2 = 2\mu m_q [1 - \delta \ln(M_\pi^2/\Lambda_\chi^2)]. \quad (10)$$

Additional contributions of order $M_\pi^2 \ln M_\pi^2$ are found to cancel. The α_0 term comes from the kinetic hairpin insertion, which is diagrammatically analogous to Fig. 1b.

3 Applications and Discussion

We wish to use our results to estimate the errors due to quenching. Baryon masses have the advantage that the leading non-analytic corrections are larger than the analytic ones by a power of M_π . (For meson masses, the enhancement is only logarithmic.) This absence of logarithms leads to the further advantage that no unknown scale Λ_χ is introduced. Thus, if we knew the quenched constants α , β *etc.*, we could use the difference between quenched and full non-analytic terms as an estimate of the effect of quenching. Of course, in practice we do not know the constants; we must either assume that they are similar to their values in the full theory or find quantities in which they cancel.

Such estimates will only be useful if the quark masses used in simulations are small enough for ChPT to be reliable. We first examine this question using the high statistics results for m_N from Ref. [5]. In Fig. 2 we plot m_N vs. M_π^2 at $\beta = 5.93$ (scaled using $a^{-1} = 1.63$) and make a “fit,” with zero degrees of freedom, to the functional form obtained from Eq. (9). We compare this to the “prediction” of our quenched calculation, using the full theory parameters from the fit of Ref. [4], together with the assumptions $\gamma = \alpha_0 = 0$. We find, in units of GeV,

$$\begin{aligned} \text{ChPT: } m_N &= 0.97 - 0.5 \frac{\delta}{.2} M_\pi + 3.4 M_\pi^2 - 1.5 M_\pi^3, \\ \text{fit: } m_N &= 0.96 - 1.0 M_\pi + 3.6 M_\pi^2 - 2.0 M_\pi^3. \end{aligned}$$

We conclude from this comparison that ChPT is doing a reasonable job. Most significant is that the negative curvature predicted by ChPT, due to the M_π^3 term, is seen with the correct magnitude. The “hook” at small m_q is due to the M_π term, which is absent in the full theory. The rough agreement on the sign and magnitude of this term is fortuitous, since both could be altered by adding in M_π^4 corrections.

Proceeding next to quantitative estimates of quenching errors in baryon masses, we address the following question: if we could calculate dimensionless ratios (e.g. m_N/f_π) in the quenched and full theories, *using the physical quark masses*,³ by how much would these ratios differ? The use of dimensionless ratios allows us to avoid worrying about overall differences in scale. First, consider the ratio m_N/f_π . The key observation here is that kaon loops do not contribute in the QA:

$$\left(\frac{m_N}{f_\pi}\right)^{(Q)} = \left(\frac{\overline{m}}{f}\right)^{(Q)} [1 + \text{pion terms} + \dots],$$

³We actually consider the slightly unphysical case of $m_u^{\text{lat}} = m_d^{\text{lat}} = (m_u^{\text{phys}} + m_d^{\text{phys}})/2$.




Figure 2: The nucleon mass from the lattice [6]. The curve is from a fit to the form $m_N = \overline{m} + a(M_\pi^2)^{1/2} + b(M_\pi^2) + c(M_\pi^2)^{3/2}$.

whereas they do in the full theory:

$$\left(\frac{m_N}{f_\pi}\right)^{(F)} = \left(\frac{\overline{m}}{f}\right)^{(F)} [1 + c_2 M_K^2 - c_3 M_K^3 + c_4 M_K^2 \ln(M_K^2/\Lambda_\chi^2) + \text{pion terms} + \dots].$$

The coefficients c_i can be obtained from the literature. The chiral expansion in the two theories is clearly quite different. To make a quantitative estimate, we ignore the pion terms (for physical quark masses, the kaon terms are dominant), we assume that $(\overline{m}/f)^{(F)} \simeq (\overline{m}/f)^{(Q)}$, and we evaluate the c_i using the fit of [4]. We find that the ratio is 16% too large in the quenched approximation. If we set the scale using f_π , this corresponds to a 150 MeV overestimate of m_N . The error could be smaller if the difference in the chiral loops were compensated by a change in $\overline{m}/f$, but this could only occur at a single value of M_K .

This estimate is about as good as one could hope to make, because only the lowest order coefficients $\overline{m}$ and f appear in the quenched result. We need not make any assumptions about the higher order coefficients $D^{(Q)}$, $F^{(Q)}$, *etc.*

We can now do a similar analysis for all the members of the baryon octet. Rather than computing the ratios to f_π , let us instead consider ratios within the octet itself, $R_{ij} \equiv m_i/m_j$. These quantities have the advantages that (a) they do not involve the unknown scale Λ_χ , and (b) if we take their ratio in the full and quenched theories, then the tree-level mass terms cancel.⁴ As above, we consider physical quark masses and therefore drop terms from pion loops, which are numerically small. This leaves contributions from loops containing kaons

⁴They cancel assuming that we ultimately equate the quenched theory parameters with their full theory counterparts, as we have done above, and which we must do to obtain quantitative estimates. Henceforth this assumption will be implied, and we drop the (Q) and (F) superscripts.

and $\bar{s}s$ -mesons, the latter type being replaced by the η in the full theory. In the QA, we find

$$\begin{aligned}
R_{\Sigma N}^{(Q)} &= 1 - \left[3D^2 - 14DF + 11F^2 \right] \delta \frac{\pi}{\sqrt{2}} \frac{M_K}{\bar{m}} + \\
&\quad - \left[-\frac{2}{3}D^2 + 4DF - 2F^2 - 4\sqrt{2}D\gamma + 4\sqrt{2}F\gamma \right] \frac{M_K^3}{16\pi f^2 \bar{m}}, \\
R_{\Xi N}^{(Q)} &= 1 - \left[-8DF + 20F^2 \right] \delta \frac{\pi}{\sqrt{2}} \frac{M_K}{\bar{m}} + \\
&\quad - \left[\left(-\frac{2}{3} - \frac{4\sqrt{2}}{3} \right) D^2 + 4DF + \left(-2 + 4\sqrt{2} \right) F^2 + 8\sqrt{2}F\gamma \right] \frac{M_K^3}{16\pi f^2 \bar{m}}, \\
R_{\Lambda N}^{(Q)} &= 1 - \left[-\frac{13}{9}D^2 - \frac{2}{3}DF + 11F^2 \right] \delta \frac{\pi}{\sqrt{2}} \frac{M_K}{\bar{m}} + \\
&\quad - \left[-\frac{10}{9}D^2 + \frac{4}{3}DF + 2F^2 + \frac{4\sqrt{2}}{3}D\gamma + 4\sqrt{2}F\gamma \right] \frac{M_K^3}{16\pi f^2 \bar{m}},
\end{aligned}$$

where we have excluded both the $O(M_K^2)$ terms, which cancel in the final ratios, and the M_K^3 terms proportional to α_0 , which vanish under our usual assumption that $\alpha_0 \simeq 0$. In the full theory, the corresponding results are [6, 4]

$$\begin{aligned}
R_{\Sigma N}^{(F)} &= 1 - \left[\left(\frac{1}{3} + \frac{4}{3\sqrt{3}} \right) D^2 + \left(2 + \frac{8}{3\sqrt{3}} \right) DF - \left(1 + \frac{4}{\sqrt{3}} \right) F^2 \right] \frac{M_K^3}{16\pi f^2 \bar{m}}, \\
R_{\Xi N}^{(F)} &= 1 - \left(4 + \frac{16}{3\sqrt{3}} \right) DF \frac{M_K^3}{16\pi f^2 \bar{m}}, \\
R_{\Lambda N}^{(F)} &= 1 - \left[\left(-1 + \frac{4}{3\sqrt{3}} \right) D^2 + \left(2 + \frac{8}{3\sqrt{3}} \right) DF + \left(3 - \frac{4}{\sqrt{3}} \right) F^2 \right] \frac{M_K^3}{16\pi f^2 \bar{m}}.
\end{aligned}$$

The ratios $r_{ij} \equiv R_{ij}^{(Q)}/R_{ij}^{(F)}$ determine the relative quenching error in the octet mass splittings. Given the assumptions stated above and taking the full-theory parameters from the first fit of Ref. [4], we obtain the following estimates:

$$\begin{aligned}
r_{\Sigma N} &= 1 + 0.19(\delta/0.2) + 0.13 = 1.31 [1.22] \text{ for } \delta = 0.2 [0.1], \\
r_{\Xi N} &= 1 - 0.46(\delta/0.2) + 0.43 = 0.97 [1.20] \text{ for } \delta = 0.2 [0.1], \\
r_{\Lambda N} &= 1 - 0.39(\delta/0.2) + 0.26 = 0.87 [1.06] \text{ for } \delta = 0.2 [0.1].
\end{aligned}$$

Clearly we need to know δ to draw quantitative conclusions. The expectation using continuum parameters is $\delta = 0.2$ [2], while the direct evidence suggests a smaller number [7]. Nevertheless, the results indicate that the pattern of splittings within the octet might differ by 20-30% in the quenched approximation from that in the full theory.

How do these estimates compare to numerical results? We can only make the comparison for m_N/f_π , since the most accurate baryon mass calculations have only been done for degenerate quarks. Ref. [5, 8] finds that this ratio is about 15% too large, in (overly) good agreement with our result. They find, however, that the ρ and decuplet baryon masses are also about 15% too large when compared to f_π , and they suggest a picture in which the baryon masses are in the correct ratio, but that f_π is too small. This hypothesis is in conflict

with our estimates for the remainder of the octet masses. It will be very interesting to see how simulations of quenched baryons with non-degenerate quarks turn out.

There is one caveat in the above comparison: Ref. [5] obtains their value for m_N using a linear extrapolation in m_q , whereas our calculation includes terms proportional to $m_q^{1/2}$ and $m_q^{3/2}$ (see Fig. 2). In other words the quenched computation of Ref. [5] is providing a quantity slightly different from $m_N^{(Q)}$ calculated in quenched CHPT. It should be straightforward to account for this difference, and we will present results in a future write-up [9].

Finally, we note that a shortcoming in the calculation discussed above is that it does not include decuplet baryons, neither as external states nor in loops. In the continuum, it has been found that the chiral expansion is, in general, better behaved if one includes the decuplet [6, 4]. This is likely to be even more true when trying to describe lattice results, because the octet-decuplet splitting in the QA is considerably smaller. It is also interesting to work out the chiral expansions for the decuplet masses, since they are computed in present simulations and allow further tests of our methodology. We have nearly completed these calculations and will present results in [9]. One result of note is that we find that the quenched Δ can decay to $p + \pi$.

Acknowledgements

We would like to thank Claude Bernard, Maarten Golterman, Paul Mackenzie, Don Weingarten, and Larry Yaffe for useful conversations. This work is supported in part by the DOE through grant DE-FG06-91ER40614, and by an Alfred P. Sloan Fellowship.

References

- [1] Claude Bernard and Maarten Golterman, Phys. Rev. D **46**, 853 (1992).
- [2] Stephen R. Sharpe, Phys. Rev. D **41**, 3233 (1990).
- [3] Elizabeth Jenkins and Aneesh V. Manohar, Phys. Lett. B **255**, 558 (1991).
- [4] V. Bernard, N. Kaiser and U. Meißner, Z. Phys. **C60**, 111 (1993).
- [5] F. Butler, H. Chen, A. Vaccarino, J. Sexton and D. Weingarten, Phys. Rev. Lett. **70**, 2849 (1993).
- [6] Elizabeth Jenkins, Nucl. Phys. B **368**, 190 (1992).
- [7] Y. Kuramashi, M. Fukugita, H. Mino, M. Okawa and A. Ukawa, these proceedings.
- [8] F. Butler, H. Chen, J. Sexton, A. Vaccarino, D. Weingarten, *Meson Decay Constants from the Valence Approximation to Lattice QCD*, IBM-HET-93-3, (Oct 1993), hep-lat/9310009
- [9] J. Labrenz and S. Sharpe, in preparation.